

## Preisträger Marathon-Preis Lauréats prix marathon

Jahr Année	Preisträger Lauréat	Titel Titre
2012	Victor Dubois-Ferrière	Predictive values of indirect detection of <i>Kingella kingae</i> osteoarticular infections in young children by PCR assays on throat's swab: toward a novel diagnostic method
	Mazda Farshad	Does subacromial injection of a local anesthetic influence strength in healthy shoulders?: a double blinded, placebo-controlled study
	Martin Clauss	Risk factors for aseptic loosening of Muller type straight stems: a register-based analysis of 828 consecutive cases with a minimum follow-up of 16 years
2011	Martin Clauss	Factors associated with rifampin resistance in staphylococcal periprosthetic joint infections (PJI): a matched case-control study
	Emin Aghayev	Benchmarking across spine registries: comparison of pain alleviation after lumbar tda and alif with stratification by surgeon
2010	Paul-Michael Bodler	Function and flexion after TKA – does design matter?
	Fabian Krause	The impact of complications in total ankle replacement and ankle fusion analyzed by prospective validated outcome at intermediate follow-up
	Jan Van Aaken	Value of sonography to detect occult fracture of the scaphoid: a comparison with CT
2009	Anne F. Mannion	The effect of post-surgical rehabilitation on outcome after surgical decompression of the lumbar spine : a randomized controlled trial with 5-year follow-up
	Andreas Krieg	Swiss multicentric study on the long-term results of patients with synovial sarcoma
	F. Unno-Veith	Clinical and radiological results 20 years after lateral meniscectomy in stable knees
2008	Domizio Suvà	Preoperative patient education reduces the incidence of dislocation after primary total hip arthroplasty
	Dimitri Ceroni	

	Frank Kleinstück	Outcome of the lower limb fractures in children and adolescents: preliminary results  A prospective study of the influence of preoperative back pain on the outcome of lumbar decompression surgery
2007	Carlo Camathias	Pin-Pflege des Fixateur externe – ein notwendiges Übel?
	Helge Burmeister	Conservative treatment of the stable lateral malleolar fracture: presentation of the outcome in a retrospective study
	Karlmeinrad Giesinger	Conventional vs. navigated total knee arthroplasty: a single surgeon comparative study
	D. Heckenbichler	Reliabilität des Verdachtbefundes “ungleiche Knochendichte” am AP-Röntgenbild der Kniegelenke im Seitenvergleich
2006	Linas Jankauskas	Loss of the subchondral sclerosis on ap radiographs of the shoulder: a pathognomonic sign for an osseous glenoid rim defect in anterior shoulder instability
	K. Tecklenburg	CT and MRI controlled prospective evaluation of new composite screws and allograft bone screws for BPTB-graft fixation in ACL-reconstruction
	Andreas Louis von Roll	Evolution of non-operatively treated supraspinatus tears
2005	Markus Kuster	A prospective randomised study to compare the lateral and medial approach for total knee arthroplasty
	R. Granes	Markraumaufbohrung als Therapie bei chronischer endomedullärer Osteomyelitis, ein 15 Jahres Follow up
	S. Blumenthal	Postoperative analgesia for anterior correction of thoracic scoliosis using double epidural catheter with continuous ropivacaine
	H. Frauenfelder-Erdogan	The association between cervical spine curvature and neck pain
2004	Arndt von Campe	Closed suction wound drainage versus non drainage in primary total hip replacement (thr). Short term difference.
	P. Szelloe	Double epidural catheter withropivacaine versus intravenous morphine for postoperataive analgesia after scoliosis correction

	L.P. Staub	Does the womac pain score deliver the information we are looking for?
2003	Urs Müller  Claudio Dora  Martin Lüem	Norm values on activities of daily living for the Swiss German population  Recurrent lumbar disc herniation: risk factors on MRI  5-J-Resultate zweier zementierter Geradschaftssysteme (Virtec. vs. Müller) in einer prospektiv randomisierten Studie
2002	Christoph Röder   Bernhard Jost  Samuel Mpyysi	Die Übereinstimmung zwischen dem Harris Hip Score und der subjektiven Patientenzufriedenheit nach totaler Hüftarthroplastik" (Co-Autoren: S. Eggli, M. Lavigne, U. Müller, A. Busato (Bern))  Die Validität der klinischen Untersuchung zur Diagnose einer Komponentenlockerung nach totaler Hüftarthroplastik" (Co-Autoren: S. Eggli, M. Aebi, A. Busato (Bern))  Klinische, funktionelle und radiologische Langzeitresultate nach totaler Hüftarthroplastik - Ergebnisse vom M.E. Müller-Hüftregister" (Co-Autoren: S. Eggli, A. Busato (Bern))  Prävalenz und Relevanz von pathologischen MRI-Befunden bei Wurfsportlern  Recommandations pour la pratique clinique (RPC) en orthopédie: quel choix de prothèse de hanche (PTH): cimentée ou non-cimentée?
2001	Ivan Broger  Peter Ochsner  Heike Bischoff	2 Jahres-Kontrollen nach Hüft- und Kniearthroplastik bei Cox- bzw. Gonarthrosepatienten mit WOMAC / SF-36- klinische Qualitätssicherung?  10 Jahre Verlaufsbeobachtung nach zementierter Schaftrevision im Vergleich zur primären Hüfttotalprothese. Klinik und Revisionsrate  Sturzprophylaxe im Alter durch Vitamin D und Kalziumzufuhr: eine randomisierte Doppelblindstudie