

## Review

# Management of Asymptomatic Bacteriuria, Urinary Catheters and Symptomatic Urinary Tract Infections in Patients Undergoing Surgery for Joint Replacement: A Position Paper of the Expert Group 'Infection' of *swissorthopaedics*

Parham Sendi<sup>1, 2</sup>✉, Olivier Borens<sup>3</sup>, Peter Wahl<sup>4</sup>, Martin Clauss<sup>5</sup>, Ilker Uçkay<sup>6, 7</sup>; for the expert group 'Infection' of *swissorthopaedics*

1. Department of Infectious Diseases, Bern University Hospital, University of Bern;
2. Institute for Infectious Diseases, University of Bern;
3. Orthopedic Septic Surgical Unit, Department of Surgery and Anaesthesiology, Lausanne University Hospital, Lausanne;
4. Division for Orthopaedic and Trauma Surgery, Cantonal Hospital Winterthur, Switzerland;
5. Clinic for Orthopedics and Trauma Surgery and Interdisciplinary Septic Surgical Unit, Kantonsspital Baselland Liestal, Switzerland;
6. Service of Infectious Diseases, Geneva University Hospitals and Faculty of Medicine, University of Geneva;
7. Orthopedic Surgery Service, Geneva University Hospitals and Faculty of Medicine, University of Geneva, Switzerland.

✉ Corresponding author: Parham Sendi, M.D., Lecturer in Infectious Diseases, Institute of Infectious Diseases, University of Bern, Switzerland. Tel +41 31 632 32 99 Fax +41 31 632 87 66 E-Mail: Parham.Sendi@ifik.unibe.ch

© Ivyspring International Publisher. This is an open access article distributed under the terms of the Creative Commons Attribution (CC BY-NC) license (<https://creativecommons.org/licenses/by-nc/4.0/>). See <http://ivyspring.com/terms> for full terms and conditions.

Received: 2017.04.05; Accepted: 2017.04.23; Published: 2017.05.25

## Abstract

In this position paper, we review definitions related to this subject and the corresponding literature. Our recommendations include the following statements. Asymptomatic bacteriuria, asymptomatic leukocyturia, urine discolouration, odd smell or positive nitrite sediments are not an indication for antimicrobial treatment. Antimicrobial treatment of asymptomatic bacteriuria does not prevent periprosthetic joint infection, but is associated with adverse events, costs and antibiotic resistance development. Urine analyses or urine cultures in asymptomatic patients undergoing orthopaedic implants should be avoided. Indwelling urinary catheters are the most frequent reason for healthcare-associated urinary tract infections and should be avoided or removed as soon as possible.

Key words: Asymptomatic bacteriuria, asymptomatic leukocyturia, urine discolouration

## Introduction

Periprosthetic joint infection (PJI) is a serious complication after arthroplasty. The relationship between abnormal results in urine analysis and potential risk for PJI often raises questions about the optimal management in patients undergoing surgery for joint replacement

In this position paper, we review definitions related to this subject and the corresponding literature. These recommendations reflect the

opinions of the expert group 'Infection' of *swissorthopaedics*, the Swiss Society of Orthopaedics and Traumatology (<http://www.swissorthopaedics.ch>).

## Definitions

Before reviewing the relation between abnormal urinary results and the risk for PJI, commonly used terms should be defined. The definitions of

bacteriuria, pyuria, leukocyturia, asymptomatic bacteriuria (ASB) and symptomatic lower urinary tract infection (UTI) are presented in **table 1** (1, 2). The method of obtaining the urine specimen matters for the definition of ASB. If the methods of obtaining urine samples are not consistent, it suggests an accuracy problem for this variable when comparing studies. Details of urinary sampling methods, the subject of complicated UTI, antibiotic treatment of symptomatic UTI, and pyelonephritis are discussed elsewhere (3), and are beyond the scope of this document. The 2014 recommendations of the Swiss Society for Infectious Diseases regarding the treatment of UTI are available on the society's website (1).

### Leukocyturia/pyuria

The differentiation between 'leukocyturia' and 'pyuria' is not common in clinical practice (**table 1**). In this document, we prefer to use the term 'leukocyturia'. It is frequently found in ASB and in other conditions associated with inflammation of the genitourinary tract (2, 4). The presence alone, or in association with nitrite positive sediments or dark-coloured urine is not reason enough to perform a urine culture or to treat an asymptomatic patient with a positive urine culture result. In other words, leukocyturia indicates an inflammation in the genitourinary tract but not necessarily an infection (low positive predictive value). In contrast, absence of leukocyturia does not rule out urinary tract infection (4-6).

### Asymptomatic bacteriuria is common

ASB is very common. Its reported prevalence is higher in older persons, in women, and in the presence of genitourinary abnormalities (2, 7). The prevalence of ASB in a general hospital population on the day of admission was recently reported to be 8.5% (8). In diabetic adults and community-dwelling elderly adults, rates of up to 50% or more, have been published (reviewed in (9)). In patients with a chronic indwelling urinary catheter, the rate of ASB is even higher (e.g., up to 100%, reviewed in (10)(11)).

### Treatment of ASB in patients undergoing surgery for joint replacement

In previous decades, various orthopaedic centres reported their practice of antimicrobial treatment for preoperatively detected ASB. In these centres, no postoperative haematogenous seeding of bacteria to the artificial joint was observed (12-14). This finding led to the rationale to screen urine preoperatively and,

upon detection of bacterial growth, to treat ASB with antibiotics in patients who were undergoing surgery for joint replacement (15, 16). In addition, surgical intervention was often delayed or, alternatively, an 8- to 10-day postoperative course of oral antibiotics was recommended (16, 17). From today's perspective and in our view, this rationale is not correct, as outlined with following data.

In 1987, a prospective study on 227 patients undergoing hip or knee replacement found no correlation between PJI and urinary tract infection (18). In 2009, Koulouvaris *et al.* examined the medical records of 19'735 patients (19). The authors found no association between preoperative or postoperative UTI and surgical site infection. The power of the study was low, mainly because the PJI rate was low (0.29%). Nonetheless, this points towards a very low risk of haematogenous seeding of microorganism from the urinary tract to the newly implanted artificial joint. This statement is supported by results from Uçkay *et al.* (11). In their retrospective study, the cumulative duration of estimated urinary tract bacterial colonization for 6'101 total joint arthroplasties patients was approximately 120'000 patient-days (median duration 295 days). In this patient population no immediate haematogenous seeding from the urinary tract to the newly implant was observed (11).

After 2013, studies performed in Spain (20, 21), in Switzerland (22), and a multinational collaboration study (Portugal, Spain, United Kingdom) (23) questioned the benefit of treating ASB. These studies included 215 (21), 471 (20), 510 (22) and 2497 (23) patients, respectively, and the results of these investigations were similar. First, in the few PJIs that occurred postoperatively, the organism that caused the infection was different from that cultured in the preoperatively obtained urine. Second, antimicrobial treatment of ASB did not effectively result in a sterile postoperative urine culture but in change of cultured organisms. For example in the Swiss study, perioperative prophylaxis with cefuroxime changed the culture results from Gram-negative to Gram-positive organisms in 50% of asymptomatic patients (22). Third, lack of antibiotic treatment for ASB was not associated with a higher risk of postoperative PJI due to a microorganism previously found in the urine or a higher risk of postoperative symptomatic UTI. These findings are in line with results from large meta-analyses showing no differences between antibiotic treatment and no treatment of ASB for the development of symptomatic UTI (24).

**Table 1.** Definitions for terms used to describe abnormal findings in urine analysis.

| Term   | Description  | Definition  | Details   |
|--|--|---|---|
| Bacteriuria  | Presence of bacteria in urine; density is expressed as colony forming units (cfu) organism per millilitre (mL)   | Presence or absence of symptoms must be determined  | Variables for the definition:<br>method of sampling<br>number of specimens<br>cfu organism /mL<br>gender  |
| Asymptomatic bacteriuria                             | Isolation of a specified quantitative count of bacteria in an appropriately collected urine specimen obtained from a person without symptoms or signs referable to urinary infection | $\geq 10^5$ cfu of the same organism/mL in 2 consecutive clean voided urine specimens (midstream) for women, or 1 specimen (midstream) for men  | If urine is obtained via catheterization, definition includes $\geq 10^2$ cfu organism/mL in 1 specimen.<br>In persons with indwelling urinary catheter, $\geq 10^3$ cfu organism/mL in 1 specimen. |
| Pyuria   | Presence of increased numbers of PMN* leukocytes in urine  | No uniformly accepted definition (e.g., $\geq 10^3$ WBC*/mL in non-centrifuged urine). Often referred to as marked leukocyturia   | Terms pyuria and leukocyturia often used interchangeably  |
| Leukocyturia   | Presence of (undifferentiated) leukocytes in urine   | $\geq 10$ leukocytes per microscopic field or positive leukocyte esterase result in a reagent strip test  | The cut-off (number of leukocytes per microscopic field) can vary between laboratories  |
| Acute uncomplicated urinary tract infection in women | A symptomatic bladder infection characterized by frequency, urgency, dysuria or suprapubic pain in a woman with a normal urinary tract   | $\geq 10^2$ cfu/mL organism in voided urine or via catheterization obtained in urine from 1 specimen. In persons with indwelling urinary catheter, $\geq 10^3$ cfu organism/mL in 1 specimen. | Absence of pain and tenderness in the costovertebral angle, no fever, no bacteraemia, no functional or structural abnormalities of the genitourinary tract  |

\* PMN, polymorphonuclear; WBC, white blood cells

The data indicates, however, that the presence of ASB prior to surgery has no clinical relevance, and therefore, should neither indicate preoperative antimicrobial treatment, nor delay surgery, nor lead to prescription of postoperative antibiotics. Also, routine perioperative antibiotic prophylaxis should not be switched to a compound active against microorganisms found in the urine of patients with ASB. Cordero-Ampuero *et al.* (20) randomly assigned patients undergoing surgery for total hip arthroplasty or hemiarthroplasty and with ASB to receive specific antibiotics ( $n = 117$ ) or routine antibiotic prophylaxis ( $n = 126$ ). Thirteen patients developed PJI after 3 months. Bacteria cultured from the wound were not those cultured in urine samples in any case, irrespective of the previous choice of antimicrobial compound.

In the work by Sousa *et al.* (23), the presence of ASB was associated with a higher risk for PJI, irrespective of organisms found in the urine. In our view, the interpretation of this finding falls into the category of weighting patients' comorbidities for postoperative complications, including infections. In other words, a wide range of patient-related factors increases the risk of surgical site infections (25, 26). Sousa *et al.* (23) and the accompanying editorial note by Duncan (27) suggested that ASB may also be a surrogate marker for increased risk of infection, but not a risk factor itself. In other words, the study suggests an epidemiological association but not a causal relation between ASB and development of a PJI of a non-urinary origin (28).

Many of the cited studies face the criticism of being underpowered. Indeed, Bouvet *et al.* (22) calculated a sample size of  $2 \times 50'979$  arthroplasties

for a superiority trial ( $\alpha 0.05$ , power 80%, difference in infection risk 0.15%) and approximately  $2 \times 200'000$  for a non-inferiority trial. It is not realistic that such a trial will ever be performed. However, Bouvet *et al.* (22) evaluated the costs associated with the screening and treatment for ASB (Geneva, Switzerland). On the basis of 510 elective arthroplasties performed in one year, the estimated minimal laboratory costs for preoperative urinary analyses were 25'000 Swiss Francs. The estimated savings by withholding antibiotics for ASB were at least 2'000 Swiss Francs (22). Understandably, this cost estimate will be likely higher when ASB with multi-drug-resistant bacteria are treated.

The power calculation is also important to highlight collateral damages caused by unnecessary use of antibiotics. Antibiotic treatment for ASB is associated with significantly more adverse events and the development of resistance (24, 29). Unfortunately, ASB has become a major cause of inappropriate antimicrobial use (30, 31). This reasoning coincides with epidemiological data showing that the rate of ASB is higher in older persons and in the presence of genitourinary abnormalities (2, 32). Evidently, these patient groups overlap with those requiring arthroplasty for osteoarthritis.

Based on these arguments, we do not recommend urine analyses in asymptomatic patients undergoing surgery for orthopaedic implants. Our recommendation is valid for all asymptomatic patients regardless of gender, age, local urologic cancer or chronic urinary catheter use. We also support this recommendation for patients treated with immunosuppressive compounds, although few data on PJI are available for this patient population.

The rationale for this recommendation is based on the observation of similar low PJI rates due to Gram-negative rods or enterococci (i.e., microorganisms typically found in the urine) in the general population (33, 34) and in a nested survey within the Swiss Transplant Cohort Study (*Ilker Uçkay, Andrey Diego, unpublished data*). Similarly, Vergidis *et al.* (35) retrospectively reviewed the arthroplasty cohort of the Mayo Clinic. Among 367 solid organ transplant patients, 12 PJIs were identified. Eight of them were caused by staphylococci and streptococci, two by nontuberculous mycobacteria, and two cases were culture-negative. Only one co-infection with *Klebsiella* spp. (0.27%) among all transplant patients was found.

If a urinary culture is nonetheless performed, a positive result should not be treated in asymptomatic patients. The routine perioperative antibiotic prophylaxis should not be modified. Moreover, the results, including the presence of leukocyturia, urine decolouration, odd smell or positive nitrite sediments should not lead to postponement of the surgical intervention (36).

### **Symptomatic bacteriuria/urinary tract infection**

The diagnosis and treatment of UTIs is discussed elsewhere (37, 38). Guideline recommendations for the treatment of UTI vary between countries and institutions on the basis of local resistance epidemiology and results from surveillance systems. The 2014 recommendations of the Swiss Society for Infectious Diseases regarding the treatment of UTI are available on the society's website (1). The decision to perform surgery after a UTI has been identified preoperatively depends on the extent of infection and the timing of surgery. In patients with signs of sepsis (e.g., urosepsis or positive blood cultures), a delay of surgery is recommended until the infection is controlled. Such a recommendation is, however, less clear for uncomplicated cystitis without sepsis. In our view, it is not necessary to postpone surgery, though antimicrobial treatment for UTI must be initiated promptly. The scientific rationale for this opinion statement however remains to be elaborated. In these cases, treatment recommendations do not differ from those for UTI in other patients (1). This is also true for postoperative UTI.

### **Prevention of postoperative urinary tract infection**

There are conflicting results about whether postoperative UTI is a risk factor for PJI. While several studies have suggested that postoperative UTI is

indeed a risk factor for PJI (26, 39), a recent review and meta-analysis did not confirm this hypothesis (40). Because the precise definitions of UTIs in these large data analyses are frequently missing, the interpretation of the results becomes ambiguous. Nonetheless, healthcare-associated infection, including symptomatic UTI after joint replacement, must be avoided.

The presence of a urinary catheter is the most important risk factor for healthcare-associated UTI (41). Indwelling urinary catheters in place for  $\geq 2$  days result in a significant increase in nosocomial infections (42). Thus, infection control strategies to prevent UTI are pivotal for every orthopaedic centre. Stéphan *et al.* (43) demonstrated that multifaceted interventions after orthopaedic surgery led to a reduction in the number of UTIs to two-thirds of that in the control group. These interventions included restricting urinary catheterization to a defined patient group, inserting catheters by using aseptic techniques and sterile equipment, connecting catheters to a closed drainage system, and removing the urinary catheter (if insertion was necessary at all) before patient discharge from the postanesthesia care unit (43). In addition, specifically tailored, locally developed guidelines were distributed; educational sessions were performed; and posters with a visual display of the guidelines were hung up. Guidelines were applied in the operating room, postanesthesia care unit and surgical ward (43). These combined efforts demonstrate the multidisciplinary need to implement prevention strategies. It is important to maintain a suitable educational level in order to sustain the high success rate of such interventions (44). This level can be achieved by engaging local leaders or a multidisciplinary team who are in decision-making positions (44).

### **Conclusions**

For ASB, a paradigm shift from treatment to non-treatment has occurred in recent years. Treating ASB does more harm than good and should be avoided. In addition, the presence of ASB should not postpone surgery. Perioperative antibiotic prophylaxis should not be modified because of ASB. This recommendation is equally valid for the presence of asymptomatic leukocyturia or positive nitrite sediments.

The most common risk factors for postoperative healthcare-associated UTIs are indwelling urinary catheters. They should be indicated for only a few patient groups and removed as rapidly as possible. Preventive strategies require a multifaceted approach and constant engagement by all involved parties.

## Acknowledgment

We thank Prof. Werner Zimmerli, MD and Dr. Andreas Kronenberg, MD for valuable discussion on the content of the manuscript. We thank the expert group for urinary tract infections of the Swiss Society of Infectious Diseases and the directive committee of Swiss Society of Infectious Diseases for endorsing this document.

Barbara Every, ELS, of BioMedical Editor, St. Albert, Alberta, Canada, provided English language editing. A German and French version of this article will be published in the journal "Swiss Medical Forum" (available from "http://www.medicalforum.ch").

## Expert group 'Infection' of swissorthopaedics

(In alphabetical order, Last Name followed by first name) Achermann Yvonne<sup>a</sup>, Borens Olivier<sup>b</sup>, Broger Ivan<sup>c</sup>, Clauss Martin<sup>d</sup>, Eich Gerhard<sup>e</sup>, Nötzli Hubert<sup>f</sup>, Nowakowski Andrej Maria<sup>g</sup>, Ochsner Peter<sup>h</sup>, Sendi Parhami<sup>i</sup>, Strahm Caroli, Suvà Domizio<sup>k</sup>, Uçkay Ilker<sup>k,l</sup>, Vogt Markus<sup>m</sup>, Wahl Peter<sup>n</sup>.

<sup>a</sup>Division of Infectious Diseases and Hospital Epidemiology, University Hospital Zurich, University of Zurich, Zurich, Switzerland.

<sup>b</sup>Orthopedic Septic Surgical Unit, Department of Surgery and Anesthesiology, Lausanne University Hospital, Lausanne, Switzerland.

<sup>c</sup>Clinic for Orthopedics and Trauma Surgery, Cantonal Hospital Graubünden, Chur, Switzerland.

<sup>d</sup>Clinic for Orthopedics and Trauma Surgery and Interdisciplinary Septic Surgical Unit, Kantonsspital Baselland Liestal, Switzerland.

<sup>e</sup>Unit of Infectious Diseases, Division of Internal Medicine, Triemlispital, Zurich, Switzerland.

<sup>f</sup>Sonnenhof Orthopaedic Center, Bern, Switzerland.

<sup>g</sup>Orthopaedic Department, University of Basel, Basel, Switzerland.

<sup>h</sup>Orthopaedic Surgeon, Professor emeritus University of Basel, Frenkendorf, Switzerland.

<sup>i</sup>Department of Infectious Diseases, Bern University Hospital, University of Bern and Institute for Infectious Diseases, Faculty of Medicine, University of Bern, Switzerland.

<sup>j</sup>Department of Infectious Diseases, Cantonal Hospital St. Gallen, St. Gallen, Switzerland.

<sup>k</sup>Orthopedic Surgery Service, Geneva University Hospitals and Faculty of Medicine, University of Geneva, Switzerland.

<sup>l</sup>Service of Infectious Diseases, Geneva University Hospitals and Faculty of Medicine, University of Geneva, Switzerland.

<sup>m</sup>Infectious Diseases Service, Cantonal Hospital Zug, Baar, Switzerland.

<sup>n</sup>Division for Orthopaedic and Trauma Surgery, Cantonal Hospital Winterthur, Switzerland.

## Competing Interests

The authors have declared that no competing interest exists.

## References

- [Internet] Hasse B, Huttner A, Huttner B, Egger M, Zanetti G, Marschall J, et al. Behandlung von unkomplizierten Harnwegsinfektionen 22 May 2014. [http://www.sginf.ch/files/behandlung\\_von\\_unkomplizierten\\_harnwegsinfektionen.pdf](http://www.sginf.ch/files/behandlung_von_unkomplizierten_harnwegsinfektionen.pdf).
- Nicolle LE, Bradley S, Colgan R, Rice JC, Schaeffer A, Hooton TM, et al. Infectious Diseases Society of America guidelines for the diagnosis and treatment of asymptomatic bacteriuria in adults. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2005;40(5):643-54.
- Nicolle LE. A practical guide to antimicrobial management of complicated urinary tract infection. *Drugs Aging*. 2001;18(4):243-54.
- Aguirre-Avalos G, Zavala-Silva ML, Diaz-Nava A, Amaya-Tapia G, Aguilar-Benavides S. Asymptomatic bacteriuria and inflammatory response to urinary tract infection of elderly ambulatory women in nursing homes. *Arch Med Res*. 1999;30(1):29-32.
- Tambyah PA, Maki DG. The relationship between pyuria and infection in patients with indwelling urinary catheters: a prospective study of 761 patients. *Arch Intern Med*. 2000;160(5):673-7.
- Ouslander JG, Schapira M, Schnelle JF, Fingold S. Pyuria among chronically incontinent but otherwise asymptomatic nursing home residents. *J Am Geriatr Soc*. 1996;44(4):420-3.
- Nicolle LE. Asymptomatic bacteriuria. *Current opinion in infectious diseases*. 2014;27(1):90-6.
- Trevino SE, Henderson JP, Wu J, Cass C, Marschall J. Prevalence of Asymptomatic Bacteriuria in Hospitalized Patients. *Infection control and hospital epidemiology : the official journal of the Society of Hospital Epidemiologists of America*. 2016;37(6):749-51.
- Ipe DS, Sundac L, Benjamin WH, Jr., Moore KH, Ulett GC. Asymptomatic bacteriuria: prevalence rates of causal microorganisms, etiology of infection in different patient populations, and recent advances in molecular detection. *FEMS Microbiol Lett*. 2013;346(1):1-10.
- Nicolle LE. The Paradigm Shift to Non-Treatment of Asymptomatic Bacteriuria. *Pathogens*. 2016;5(2).
- Uckay I, Lubbeke A, Emonet S, Tovmirzaeva L, Stern R, Ferry T, et al. Low incidence of haematogenous seeding to total hip and knee prostheses in patients with remote infections. *The Journal of infection*. 2009;59(5):337-45.
- Donovan TL, Gordon RO, Nagel DA. Urinary infections in total hip arthroplasty. Influences of prophylactic cephalosporins and catheterization. *The Journal of bone and joint surgery American volume*. 1976;58(8):1134-7.
- Glynn MK, Sheehan JM. The significance of asymptomatic bacteriuria in patients undergoing hip/knee arthroplasty. *Clinical orthopaedics and related research*. 1984;185:151-4.
- Kumar P, Mannan K, Chowdhury AM, Kong KC, Pati J. Urinary retention and the role of indwelling catheterization following total knee arthroplasty. *Int Braz J Urol*. 2006;32(1):31-4.
- Singh D, Roberts C, Bentley G. Urinalysis before joint arthroplasty. To dipstick or not? That is the question. *Ann R Coll Surg Engl*. 1998;80(4):300.
- Rajamanickam A, Noor S, Usmani A. Should an asymptomatic patient with an abnormal urinalysis (bacteriuria or pyuria) be treated with antibiotics prior to major joint replacement surgery? *Cleveland Clinic journal of medicine*. 2007;74 Suppl 1:S17-8.
- David TS, Vrahas MS. Perioperative lower urinary tract infections and deep sepsis in patients undergoing total joint arthroplasty. *J Am Acad Orthop Surg*. 2000;8(1):66-74.
- Ritter MA, Fechtman RW. Urinary tract sequelae: possible influence on joint infections following total joint replacement. *Orthopedics*. 1987;10(3):467-9.
- Koulouvaris P, Sculco P, Finerty E, Sculco T, Sharrock NE. Relationship between perioperative urinary tract infection and deep infection after joint arthroplasty. *Clin Orthop Relat Res*. 2009;467(7):1859-67.
- Cordero-Ampuero J, Gonzalez-Fernandez E, Martinez-Velez D, Esteban J. Are antibiotics necessary in hip arthroplasty with asymptomatic bacteriuria? Seeding risk with/without treatment. *Clin Orthop Relat Res*. 2013;471(12):3822-9.
- Martinez-Velez D, Gonzalez-Fernandez E, Esteban J, Cordero-Ampuero J. Prevalence of asymptomatic bacteriuria in knee arthroplasty patients and subsequent risk of prosthesis infection. *Eur J Orthop Surg Traumatol*. 2016;26(2):209-14.
- Bouvet C, Lubbeke A, Bandi C, Pagani L, Stern R, Hoffmeyer P, et al. Is there any benefit in pre-operative urinary analysis before elective total joint replacement? *Bone Joint J*. 2014;96B(3):390-4.
- Sousa R, Munoz-Mahamud E, Quayle J, Dias da Costa L, Casals C, Scott P, et al. Is asymptomatic bacteriuria a risk factor for prosthetic joint infection?

- Clinical infectious diseases : an official publication of the Infectious Diseases Society of America. 2014;59(1):41-7.
24. Zalmanovici Trestioreanu A, Lador A, Sauerbrun-Cutler MT, Leibovici L. Antibiotics for asymptomatic bacteriuria. The Cochrane database of systematic reviews. 2015;4:CD009534.
  25. Cheadle WG. Risk factors for surgical site infection. *Surgical infections*. 2006;7 Suppl 1:S7-11.
  26. Pulido L, Ghanem E, Joshi A, Purtill JJ, Parvizi J. Periprosthetic joint infection: the incidence, timing, and predisposing factors. *Clinical orthopaedics and related research*. 2008;466(7):1710-5.
  27. Duncan RA. Prosthetic joint replacement: should orthopedists check urine because it's there? *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2014;59(1):48-50.
  28. Uckay I, Lubbeke A, Huttner B. Preoperative asymptomatic bacteriuria and subsequent prosthetic joint infection: lack of a causal relation. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2014;59(10):1506-7.
  29. Cai T, Nesi G, Mazzoli S, Meacci F, Lanzafame P, Caciagli P, et al. Asymptomatic bacteriuria treatment is associated with a higher prevalence of antibiotic resistant strains in women with urinary tract infections. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2015;61(11):1655-61.
  30. Silver SA, Baillie L, Simor AE. Positive urine cultures: A major cause of inappropriate antimicrobial use in hospitals? *Can J Infect Dis Med Microbiol*. 2009;20(4):107-11.
  31. Trautner BW. Asymptomatic bacteriuria: when the treatment is worse than the disease. *Nat Rev Urol*. 2012;9(2):85-93.
  32. Nicolle LE. Asymptomatic bacteriuria in the elderly. *Infectious disease clinics of North America*. 1997;11(3):647-62.
  33. Uckay I, Bernard L. Gram-negative versus gram-positive prosthetic joint infections. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2010;50(5):795.
  34. Jamei O, Gjoni S, Zenelaj B, Kressmann B, Belaieff W, Hannouche D, et al. Which Orthopaedic Patients Are Infected with Gram-negative Non-fermenting Rods? *J Bone Joint Infect*. 2017;2(2): 73-6.
  35. Vergidis P, Lesnick TG, Kremers WK, Razonable RR. Prosthetic joint infection in solid organ transplant recipients: a retrospective case-control study. *Transpl Infect Dis*. 2012;14(4):380-6.
  36. Gou W, Chen J, Jia Y, Wang Y. Preoperative asymptomatic leucocyturia and early prosthetic joint infections in patients undergoing joint arthroplasty. *J Arthroplasty*. 2014;29(3):473-6.
  37. Gupta K, Hooton TM, Naber KG, Wullt B, Colgan R, Miller LG, et al. International clinical practice guidelines for the treatment of acute uncomplicated cystitis and pyelonephritis in women: A 2010 update by the Infectious Diseases Society of America and the European Society for Microbiology and Infectious Diseases. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2011;52(5):e103-20.
  38. Hooton TM, Bradley SF, Cardenas DD, Colgan R, Geerlings SE, Rice JC, et al. Diagnosis, prevention, and treatment of catheter-associated urinary tract infection in adults: 2009 International Clinical Practice Guidelines from the Infectious Diseases Society of America. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2010;50(5):625-63.
  39. Wymenga AB, van Horn JR, Theeuwes A, Muijtjens HL, Slooff TJ. Perioperative factors associated with septic arthritis after arthroplasty. Prospective multicenter study of 362 knee and 2,651 hip operations. *Acta Orthop Scand*. 1992;63(6):665-71.
  40. Zhu Y, Zhang F, Chen W, Liu S, Zhang Q, Zhang Y. Risk factors for periprosthetic joint infection after total joint arthroplasty: a systematic review and meta-analysis. *J Hosp Infect*. 2015;89(2):82-9.
  41. Uckay I, Sax H, Gayet-Ageron A, Ruef C, Muhlemann K, Troillet N, et al. High proportion of healthcare-associated urinary tract infection in the absence of prior exposure to urinary catheter: a cross-sectional study. *Antimicrob Resist Infect Control*. 2013;2(1):5.
  42. Wald HL, Ma A, Bratzler DW, Kramer AM. Indwelling urinary catheter use in the postoperative period: analysis of the national surgical infection prevention project data. *Archives of surgery*. 2008;143(6):551-7.
  43. Stephan F, Sax H, Wachsmuth M, Hoffmeyer P, Clergue F, Pittet D. Reduction of urinary tract infection and antibiotic use after surgery: a controlled, prospective, before-after intervention study. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2006;42(11):1544-51.
  44. Sax H, Kuster SP, Tehrani YA, Ren R, Uckay I, Agostinho A, et al. Eight-year sustainability of a successful intervention to prevent urinary tract infection: A mixed-methods study. *Am J Infect Control*. 2016;44(7):820-4.